

Lost Terranes of Zealandia: possible development of late Paleozoic and early Mesozoic sedimentary basins at the southwest Pacific margin of Gondwanaland, and their destination as terranes in southern South America

Christopher J. Adams

GNS Science, Private Bag 1930, Dunedin, New Zealand.
argon@gns.cri.nz

ABSTRACT. Latest Precambrian to Ordovician metasedimentary successions and Cambrian-Ordovician and Devonian-Carboniferous granitoids form the major part of the basement of southern Zealandia and adjacent sectors of Antarctica and southeast Australia. Uplift/cooling ages of these rocks, and local Devonian shallow-water cover sequences suggest that final consolidation of the basement occurred through Late Paleozoic time. A necessary consequence of this process would have been contemporaneous erosion and the substantial development of marine sedimentary basins at the Pacific margin of Zealandia. These are found nowhere at the present day, suggesting that the basins have been lost by tectonic erosion, perhaps in a margin-parallel dextral translation similar to late Paleozoic-Mesozoic suspect terranes of New Zealand. A probable detrital zircon age pattern is assembled for these lost Zealandia sediments, and then compared with those of pre-Jurassic (probable Triassic to Devonian) metasedimentary rocks in the Chilean archipelago. Significant Mesoproterozoic, latest Neoproterozoic-Cambrian and Devonian-Carboniferous detrital zircon age components are common to both, thus supporting a possible Chilean terrane destination for these 'lost terranes of Zealandia'.

Keywords: Zealandia, Detrital zircon ages, Paleozoic, Terranes, Chile.

RESUMEN. Terrenos perdidos de Zealandia: posible desarrollo de cuencas sedimentarias del Paleozoico tardío y Mesozoico temprano en el margen suroccidental del Pacífico de Gondwana y su destino como terrenos en el sur de América del Sur. Las sucesiones metasedimentarias del Precámbrico tardío al Ordovícico y granitoides del Cámbrico-Ordovícico y Devónico-Carbonífero constituyen la mayor parte del basamento del sur de Zealandia y sectores adyacentes de la Antártica y el sudeste de Australia. Las edades de enfriamiento/alzamiento de estas rocas y la cobertura local de secuencias de aguas someras del Devónico, sugieren que la consolidación definitiva del basamento se produjo durante el Paleozoico tardío. Una consecuencia necesaria de este proceso habría sido la erosión contemporánea y el desarrollo sustancial de cuencas sedimentarias marinas en el margen del Pacífico de Zealandia. Estas no se encuentran en ninguna parte en la actualidad, lo que sugiere que las cuencas se han perdido por erosión tectónica, tal vez en una traslación dextral paralela al margen similar a los terrenos del Paleozoico tardío-Mesozoico de Nueva Zelanda. Hay una agrupación que da un probable patrón de edad de circones detríticos para estos sedimentos de la perdida Zealandia, comparables con aquellos de rocas metasedimentarias del pre-Jurásico (probable Triásico a Devónico) del archipiélago chileno. Importantes componentes de edades de circones detríticos del Mesoproterozoicos, Neoproterozoico tardío-Cámbrico y Devónico-Carbonífero son comunes a ambas regiones, favoreciendo así un posible destino chileno para estos 'terrenos perdidos de Zealandia'.

Palabras clave: Zealandia, Edades de circones detríticos, Paleozoico, Terrenos, Chile.

1. Introduction

Zealandia is that part of the greater New Zealand region underlain by continental crust (Sutherland, 1999) and which formerly constituted a sector of the Pacific margin of the Gondwanaland supercontinent. A fit of Zealandia with the adjacent continental components of Gondwanaland can be made against eastern Australia by closing the Tasman Sea, and the Ross Sea regions of Antarctica by closing the southwest Pacific Ocean. This then reveals the original continuity of several late Precambrian and early Paleozoic fold belts as demonstrated for example by common sedimentary basins, plutonic arcs and metamorphic belts (Adams, 1981; Bradshaw *et al.*, 1983; Glen 2005). Despite a large proportion (95%) of Zealandia being below sea-level and unavailable for study, its geological history compares well with its Australian and Antarctic continental neighbours, but an important difference is that Silurian and Devonian sedimentary rocks are very local and limited and, apart from a single, tiny (100 m²) seamount limestone occurrence (Jenkins and Jenkins, 1971), the Carboniferous is absent.

This paper examines this striking paucity of Silurian, Devonian and Carboniferous sedimentary rocks, and proposes that they were in fact developed in quite extensive sedimentary basins along the Zealandia Pacific margin, but were then translated, as suspect terranes, to southern South America.

2. The Zealandia continental basement: a geological outline

The pre-100 Ma basement of the Zealandia continent has three main components:

- An inboard early Paleozoic, Western Province comprises, in the Buller Terrane, extensive sandstone-dominated, turbiditic rocks of Ordovician age and minor Devonian, shallow-water successions and, in the Takaka Terrane, Cambrian to Devonian, shallow-water limestones and quartzites (Cooper and Tulloch, 1992). The province probably continues southwards on to the Campbell Plateau (Adams, 2007) and into Marie Byrd Land, West Antarctica (Bradshaw *et al.*, 1997; Pankhurst *et al.*, 1998), and also westwards into the early Paleozoic, Lachlan Fold Belt of southeast Australia (Foster and Gray, 2000; VandenBerg *et al.*, 2000; Veevers *et al.*, 2000), and more distantly, to the Late

Precambrian-Cambrian, Ross and Delamerian Fold Belts of Antarctica and Australia respectively (Tessensohn *et al.*, 1981; Adams, 1997; Adams and Ireland, 2007). Undoubted Precambrian basement intrusive igneous rocks within this region are very rare (Fioretti *et al.*, 2005), but Cambrian-Ordovician granitoid complexes within the Ross/Delamerian Fold Belts are extensive in Northern Victoria Land (Tessensohn *et al.*, 1981) and South Australia. Devonian-Early Carboniferous granitoids occur in Northern Victoria Land (Tessensohn *et al.*, 1981; Tonarini and Rocchi, 1994), Marie Byrd Land (Adams, 1987), Zealandia (Muir *et al.*, 1996) and southeast Australia (Birch, 2008). Regional deformation and metamorphism throughout is late Ordovician-Silurian (Adams *et al.*, 1975; Adams and Kreuzer, 1984; Kreuzer *et al.*, 1987; Adams, 1986; Foster and Gray, 2000). In general, the fold belts represent an outward accretion in late Neoproterozoic to Ordovician times of Ross-Delamerian marginal basins in the Transantarctic Mountains and South Australia. These basins derived their sediment inputs from the immediately adjacent Australian and Antarctic Precambrian cratons. Subsequently, in southeast Australia, Zealandia and West Antarctica, Ordovician sandstone-dominated successions of the Lachlan Fold Belt (and equivalents) developed very extensively (5,000 km) at the contemporary margin of Gondwanaland (Ireland *et al.*, 1998), with reworking from both the Australian-Antarctic Precambrian cratons and the Ross-Delamerian Fold Belts. Although all these successions appear autochthonous with respect to their proposed sediment sources inboard, to harmonise their geological history in a spatial sense, Glen (2005) has nonetheless proposed major strike-slip displacement of several early Paleozoic tectonic blocks within southeast Australia and the adjacent Ross Sea regions of Antarctica.

- An outboard, Eastern Province of Late Paleozoic-Mesozoic, tectonostratigraphic terranes (Fig. 1), several of which have been regarded as allochthonous (Bishop *et al.*, 1985). Within this province there is a western terrane group represented by island-arc, volcanoclastic-dominated (mainly acid-intermediate), forearc sedimentary basins of the Brook Street, Murihiku and Dun Mountain-Maitai Terranes (Coombs *et al.*, 1976; Ballance and Campbell 1993; Landis *et al.*, 1999; Rattenbury *et al.*, 1998). Volcanic successions are




FIG. 1. The Pacific margin of Gondwanaland in a Late Triassic reconstruction showing: **left diagram:** late Neoproterozoic-early Paleozoic basement outcrop (dark grey shading) and possible geographic continuations (light grey shading) and the Precambrian craton margin (bold broken line). The bold dotted ellipses represent probable areas of late Paleozoic-Mesozoic sediment depocentres, that might now form lost terranes of Zealandia, destined for Chile (suggested tectonic transport direction shown as bold arrows); **right diagram:** late Paleozoic-Mesozoic outcrop (dark grey shading) and possible geographic continuations (light grey shading). Areas of Jurassic-Cretaceous plutonism are shown as stipple. The bold dashed ellipses represent probable areas of late Paleozoic-Mesozoic sediment depocentres that now form suspect terranes in the Eastern Province, New Zealand (bold dotted envelopes).

well represented in the Permian, Brook Street Terrane, but are rare throughout the Late Permian to Late Jurassic, Murihiku Terrane. The Dun Mountain-Maitai Terrane is dominated by the Early Permian, Dun Mountain Ophiolite Belt (serpentinites, spilites and ultramafites) (Coombs

et al., 1976), which is overlain by Permian-Early Triassic turbidite successions of the Maitai Group. An eastern terrane group, comprising the Torlesse (composite), Waipapa and Caples Terranes (Mackinnon, 1983; Adams *et al.*, 2009), forms accretionary complexes (Permian to Cretaceous)

with extensive sandstone-dominated turbiditic rocks mainly derived from an adjacent continental margin including plutonic and metamorphic rocks. Acid-intermediate, volcanoclastic sediment inputs are pervasive in the Caples Terrane, less so in the Waipapa Terrane and rare in the Torlesse Terrane. Volcanic horizons are rare in all three terranes. Regional deformation and low-grade metamorphism within the developing accretionary wedge are recorded throughout the Triassic, Jurassic and (mainly) Cretaceous (Adams *et al.*, 1985; Adams, 2003; Adams and Maas, 2004).

- A Late Permian-Early Cretaceous, Median Batholith (Kimbrough *et al.*, 1994; Mortimer *et al.*, 1999), which separates the Western and Eastern Provinces.

In the Eastern Province, the Torlesse Terrane rocks in particular show petrographic characteristics (Mackinnon, 1983), geochemical compositions (Roser and Korsch, 1999) and radiogenic isotope patterns (Pickett and Wasserburg, 1989; Frost and Coombs, 1989; Wandres *et al.* (2004a, b); Adams *et al.*, 2005) that do not match with possible sources at the immediate Gondwanaland margin in the Western Province (Adams, 1997, 2007; Gray, 1990; Adams, 2004). Furthermore, whilst they lend support to some alternative provenances, for example within the northeast Australia or Antarctic sectors, these features do not unequivocally confirm such proposals. Attention has therefore been concentrated instead upon their detrital mica and zircon age populations. These similarly exclude any likely source areas at the adjacent margin of the Western Province of New Zealand (which are mainly Cambrian-Devonian sedimentary rocks, and Devonian-Carboniferous plutonic rocks) or their equivalents in Australia and Antarctica (Ireland, 1992a; Ireland *et al.*, 1994; Adams and Kelley, 1998; Pickard *et al.*, 2000; Cawood *et al.*, 2002; Wandres *et al.*, 2004a; Adams *et al.*, 2007) (Fig. 1). However, there is disagreement about any alternative provenances. For example, Cawood *et al.* (2002) suggested that Permian sandstones in the Akatarawa microterranes (within the Torlesse Terrane) had a distant Tethyan origin. On a comparison of petrographic, geochemical, and age information, Wandres *et al.* (2004a) suggested a West Antarctic (Amundsen Sea) source for Cambrian boulders in Torlesse conglomerates. However, these source regions remain unsatisfactory, because rocks of the appropriate extent (>100 km scale), composition (acid-intermediate), and age (Permian-Triassic) are

only very locally developed (km-scale) or absent in these regions. Except for some acid volcanic rocks in Tasmania, Cambrian (and late Precambrian) igneous rocks are similarly rare or absent. No rocks of this age are known on the Campbell Plateau. Also it cannot be satisfactorily explained how Cambrian igneous rocks, occurring as clasts in rare Torlesse conglomerates (<0.01% total) have been transported large distances (>1,000 km) from tiny occurrences in West Antarctica (Amundsen Sea), whilst there is no evidence of sand-size detritus travelling similar distances from the widespread Devonian granitoids in this same sector of the Gondwanaland margin.

Alternatively, on the basis of detrital U-Pb zircon and $^{40}\text{Ar}/^{39}\text{Ar}$ mica age patterns, Adams and Kelley (1998); Pickard *et al.* (2000) and Adams *et al.* (2007) preferred Torlesse (and Waipapa) terrane sediment provenances within the Carboniferous-Triassic plutonic and volcanic complexes of the New England Fold Belt of northeast Australia, and to a lesser degree their hinterland consisting of Precambrian-early Paleozoic rocks within the Thomson and Delamerian fold belts (Veevers, 2000; Glen, 2005). A similar preference was given for rocks within the Caples Terrane (Adams *et al.*, 2009). Features such as euhedral rather than rounded zircon mineral grains (Pickard *et al.*, 2000), conglomerate horizons (Andrews *et al.*, 1976; Smale, 1980; Wandres *et al.*, 2004b), gritstones and coarse estuarine sandstones (Andrews, 1974; Retallack, 1983), and plant material (Retallack and Ryburn, 1982) all suggest that the sandstone successions formed relatively close to a contemporary active continental margin, and the resultant sedimentary basins were subsequently transported tectonically, as suspect terranes, along the Gondwanaland margin southwards to their present New Zealand position (Adams *et al.*, 2007) (Fig. 1). This model raises the possibility that equivalent rocks were originally present at the Zealandia margin and also underwent similar tectonic transport elsewhere.

3. Late Paleozoic uplift and erosion in southern Zealandia

Throughout Zealandia and neighbouring southeast Australia and Antarctica, the K-Ar and $^{40}\text{Ar}/^{39}\text{Ar}$ cooling age patterns of the early Paleozoic meta-sedimentary terranes and early and late Paleozoic plutonic complexes are all in the range 440 to ca. 340 Ma (Adams, 1981, 1986, 1987; Adams and Kreuzer, 1984; Adams *et al.*, 1975, 1979; Foster

and Gray, 2000; Kreuzer *et al.*, 1987; Vita *et al.*, 1991; Wright *et al.*, 1991; Tonarini and Rocchi, 1994). These data suggest that substantial basement uplift and erosion continued from Silurian to at least mid-Carboniferous times. This could explain the absence of Carboniferous clastic sedimentary rocks in New Zealand.

Such exhumation must have been considerable (5-15 km), to bring greenschist facies metasediments and granitoids to the surface by the Devonian in the Buller Terrane of New Zealand (Bradshaw, 1995), in Northern Victoria Land (Tessensohn *et al.*, 1981), Marie Byrd Land, Antarctica (Grindley and Mildenhall, 1981), and in Tasmania and Victoria, Australia (VandenBerg *et al.*, 2000; Birch, 2008), but later, by Late Permian time, in the Takaka Terrane of New Zealand (Rattenbury *et al.*, 1998; Bradshaw, 2000). Throughout this broad region, Devonian sedimentary cover rocks are invariably local (km-scale), and indicate prevailing shallow-water, passive margin environments. Devonian granitoids were also emplaced in all these areas too, and minor associated volcanics are recorded in Antarctica (Northern Victoria Land, and Marie Byrd Land). Thus, throughout late Paleozoic time, the Zealandia region was relatively stable, with restricted shallow-water sedimentation, and undergoing steady regional uplift.

The substantial exhumation of such a large basement region (1,000 km-scale) necessarily implies commensurate erosion and development of sedimentary basins along the contemporary Zealandia continental margin. Surprisingly, no such basins have ever been discovered. It must be concluded that they have either been completely removed by Mesozoic subaerial erosion, or, like the Eastern Province sedimentary basins, they have undergone some form of 'tectonic' erosion, and as 'Lost Terranes of Zealandia', have been transported elsewhere.

4. Sediment inputs into postulated basins at the Zealandia margin

Any late Paleozoic basins developed at the Zealandia margin would have derived their sediment inputs from an exposed basement hinterland that included:

1. Medium-high-grade metamorphic rocks (from late Neoproterozoic to Cambrian protoliths).
2. Cambrian-Ordovician granitoid complexes of the Ross and Delamerian Fold Belts.
3. Low-medium-grade Late Cambrian-Ordovician metasediments of the Lachlan Fold Belt and correlatives in Zealandia and Antarctica.
4. Devonian to Carboniferous granitoid complexes, also within the Lachlan Fold Belt and its correlatives, *e.g.*, Karamea Batholith of New Zealand.
5. Late Permian igneous rocks of the Median Batholith or sedimentary rocks of the Brook Street and Murihiku Terranes (Fig. 1).

Today, the crucial continental basement in southern Zealandia (Campbell Plateau) and West Antarctica (Marie Byrd Land) is hidden beneath sea and ice. Since proven outcrop areas are extremely limited, estimates of the extent of sources (1)-(5) are very uncertain. Bradshaw *et al.* (1997) speculated that Permian-Jurassic rocks of source (5) did indeed continue, as terranes, across the Campbell Plateau into Marie Byrd Land. In contrast, Adams (2007), citing uniform regional magnetic anomaly patterns (Sutherland, 1999), suggested the basement in this region comprised almost entirely pre-Permian sources (3) and (4), perhaps >70%.

The very extensive sandstone-dominated successions of sources (1) and (3) have plutonic, mostly granitoid, provenances, and thus, when reworked into contemporary late Paleozoic sediments, would contribute these characteristics. To this would be added similar detritus from granitoid sources (2) and (4). Thus, late Paleozoic sediments would be dominated by sands and silts with relatively quartzose compositions, more potassic than calcic, and with elevated initial Sr isotopic ratios, >0.708 (Gray, 1990; Adams, 1997). There is, however, a complication closer to the Zealandia margin (and present-day New Zealand) where Late Permian sediment contributions from source (5), such as the Median Batholith, and Brook Street and Dun Mountain-Maitai Terranes, might become important. The extension of none of these on to the Campbell Plateau is proven, although a linear magnetic anomaly extending 400 km southeast of New Zealand might be related to the Dun Mountain Ophiolite Belt (Sutherland 1999; Mortimer *et al.*, 2002). If source (5) was originally far larger, *i.e.*, a large proportion has been removed by erosion, or if substantial extensions are in fact hidden on the Campbell Plateau and in Marie Byrd Land (Bradshaw *et al.*, 1997), then sediments contributed from this source would be less quartzose,

more volcanoclastic and with lower bulk initial $^{87}\text{Sr}/^{86}\text{Sr}$ values <0.706 .

5. Detrital zircon age patterns in late Paleozoic sediments at the Zealandia margin

The sediment sources (1)-(5) would deliver detrital zircon age characteristics to the Zealandia margin as follows:

Source (1), late Neoproterozoic-Cambrian meta-sedimentary rocks, having two main, and roughly equal, groups: late Mesoproterozoic, *ca.* 1,000-1,100 Ma, and late Neoproterozoic to Cambrian, *ca.* 600-500 Ma.

Source (2), Ross-Delamerian granitoids having primary zircons, *ca.* 460-500 Ma.

Source (3), early Paleozoic metasedimentary rocks of the Lachlan Fold Belt (and its continuations) having a combined age signature made up from sources (1) and (2).

Source (4), Devonian-Carboniferous, granitoids and/or associated volcanic rocks, having primary zircons, 430-330 Ma.

Source (5), granodiorites and diorites of the Median Batholith (Mortimer *et al.*, 1999) or older, intermediate volcanoclastic rocks in the Brook Street Terrane (Landis *et al.*, 1999), Murihiku Terrane (Ballance and Campbell, 1993) or Dun Mountain-Maitai Terrane, having mostly Late Permian (260-250 Ma) zircons.

An estimate of their relative extent would suggest that sources (1)-(3) would be major zircon contributors to any Zealandia sedimentary basins, perhaps comprising 60-70% of total, with an important but minor contribution, perhaps 20-30%, from source (4), and an uncertain, but smallest input from source (5). This contrasts markedly with zircons delivered to the northeastern Australian Gondwanaland margin to form the Torlesse, Waipapa and Caples sedimentary basins, which were dominated by source (5), often 40-50%, with only a minor contribution, 30-40%, from sources (1)-(3), and very little from source (4). These patterns are exemplified in figure 2 (upper) by zircon age distributions in Rakaia Terrane rocks of the Torlesse Composite Terrane.

6. Lost terranes of Zealandia in Chilean Patagonia

If the postulated late Paleozoic sedimentary basins at the Zealandia margin did undergo a mar-

gin-parallel translation (*ca.* 2,000 km) as suspect terranes, similar to that of the Eastern Province terranes, then possible destination terranes might be represented in the Chilean archipelago of southern South America, as the Chonos, Eastern Andes, and Duque de York Metamorphic Complexes (Faúndez *et al.*, 2002; Hervé and Fanning, 2003; Hervé *et al.*, 2003; Lacassie *et al.*, 2006). These comprise poorly fossiliferous or unfossiliferous, low-grade siliciclastic metasediments, with exception of the Duque de York Complex, which has significant Permian limestones. In general, their age range is only poorly estimated as post-Early Permian, pre-Early Cretaceous. Faúndez *et al.* (2002) and Lacassie *et al.* (2006) compared Permian (and possible Triassic) rocks in Chilean Patagonia and southern New Zealand, and they noted that the Eastern Province terranes also have poorly fossiliferous low-grade, metasediments (post-Early Permian to pre-Late Cretaceous) which include important Late Permian limestones in the Brook Street and Dun Mountain-Maitai Terranes (Rattenbury *et al.*, 1998; Landis *et al.*, 1999).

Zircon age data from the Chilean complexes cited above are summarised in figure 2 (lower) in a similar manner to the equivalent data from the Permian-Triassic Rakaia Terrane of New Zealand displayed in figure 2 (upper). The Chilean zircon age patterns are seen to contrast with those from New Zealand. Since the Rakaia Terrane is allochthonous with respect to the southern Zealandia margin, it has zircon age data that do not match the possible basement sources there. However, the Chilean zircon data do conform to that expected of a Zealandia margin sedimentary basin. In particular there is a large zircon component (early Paleozoic and Precambrian) compatible with source-types (1)-(3), and a significant minor component (Devonian and Carboniferous) compatible with source-type (4). Permian zircon contributions of source-type (5) are more significant than predicted, but are highly variable, 10-70%, suggesting that their influence is more local, and perhaps related to volcanic sources.

7. Discussion

With the exception of the Duque de York Complex discussed below, Hervé and Fanning (2003) and Hervé *et al.* (2003) inferred from detrital zircon age data in the metamorphic complexes of the Chilean

DETRITAL ZIRCON AGE DATA FOR NEW ZEALAND EASTERN PROVINCE TERRANES: PERCENTAGE PROPORTIONS IN GEOLOGICAL PERIODS

TORLESSE COMPOSITE TERRANE			Cret.	Juras.	Trias.	Per.	Carb.	Dev-Sil	Ord-Cam	Precam		
Rakaia Terrane												
4	RBW1	205	220			33	19	6	2	16	24	RBW1
2	OTQ1	220	230			30	16	9	4	13	27	OTQ1
2	HERM2	220	230			29	18	5	5	15	29	HERM2
4	NGQ2	220	230	1		32	19	4	1	14	29	NGQ2
4	NGQ2	220	230			39	10	4	1	20	26	NGQ2
4	PUD1	220	230			33	23	7	1	12	23	PUD1
5	Balmacaan	235	242			14	31	9	5	11	30	Balmacaan
9	SEL4	251	260			5	55	7	5	15	13	SEL4
4	PAR2	251	260			3	68	1	6	10	13	PAR2
1	Aviemore					4	40	4		16	36	Aviemore
5	KUROW	251	260			7	38	25	7	13	10	KUROW
6	BMQX35					2	20	21	16	24	16	BMQX35
Kakahu Terrane												
4	KAK35	251				66	14	7			14	KAK35
Akatarawa Terrane												
4	TAKA10	251		6		18	19	22	7	7	28	TAKA10
3	te94	251			4	27	19	22	4		22	te94

DETRITAL ZIRCON AGE DATA FOR METAMORPHIC COMPLEXES OF CHILEAN PATAGONIA: PERCENTAGE PROPORTIONS IN GEOLOGICAL PERIODS

Data Source	Sample Name/No.	Location	Strat. range (Ma)		Zircon percentages in selected age ranges						Sample Name/No.		
			young	old	Cret.	Juras.	Trias.	Per.	Carb.	Dev-Sil		Ord-Cam	Precam
CHONOS METAMORPHIC COMPLEX													
7	CE96-03				2	16	16	8	18	24	16	CE96-03	
7	FO96-06					12	42	15	12	5	15	FO96-06	
7	CE96-25					12	20	28	15	4	22	CE96-25	
7	FO96-40							5	4	16	75	FO96-40	
EASTERN ANDES METAMORPHIC COMPLEX													
7	PE99-32A					35	5	15	5	15	15	19	PE99-32A
7	FO98-P17					2	20	17	13	22	27	FO98-P17	
7	FF99-01					4	10	8	12	10	56	FF99-01	
7	FF99-05				4		22	16	12	24	27	FF99-05	
7	S199-28C					2	6		12	24	56	S199-28C	
7	SE99-08					2	18	2	10	40	28	SE99-08	
7	VS11A							2	2	21	73	VS11A	
7	PS98-01							2		12	86	PS98-01	
MAIN RANGE METAMORPHIC COMPLEX													
7	CE96-29				2	46	10		20	10	12	CE96-29	
7	MA21								12	29	59	MA21	
DUQUE DE YORK METAMORPHIC COMPLEX													
	AL1					63	18	9	11	5	13	AL1	
8	MD3				4	68	12		8	2	6	MD3	
8	MD32					59	11		15	2	13	MD32	

References

- Ireland, 1992b
- Adams *et al.*, 1998
- Cawood *et al.*, 2002
- Pickard *et al.*, 2000
- Wandres *et al.*, 2004a
- Adams *et al.*, 2007
- Hervé *et al.*, 2003
- Herve and Fanning, 2003

Percentage abundance categories

accessory	1-4%
minor	5-19%
major	20-49%
large	50-79%
dominant	80-100%

FIG. 2. ²⁰⁶Pb/²³⁸U detrital zircon ages for late Paleozoic-Mesozoic metasedimentary rocks of the Torlesse Composite Terrane in New Zealand (upper), compiled from Adams *et al.* (2007), and metamorphic complexes in Patagonian Chile (lower), compiled from Hervé and Fanning (2003), Hervé *et al.* (2003). Age data are divided into selected geological periods, and categorised as percentages of the total zircon set.

archipelago, autochthonous, passive-margin settings for the original sedimentary basins (see also Faúndez *et al.*, 2002). In this respect, Augustsson *et al.* (2006) reached a similar conclusion for the provenance of detrital zircons in higher-grade metasediments, of probable original Carboniferous age, in the mainland part of Chilean Patagonia. Considering only the magmatic zircons in their datasets (of which age subsets are consequently often rather small), significant proportions are Mesoproterozoic (>10%), early Paleozoic (>10%), and Devonian-Carboniferous (>15%) in age, all of which are compatible, in reasonable combinations, with a Zealandia origin. However, it must be conceded that detrital zircon age patterns in comparable late Paleozoic rocks having similar Mesoproterozoic, early Paleozoic and Devonian-Carboniferous components, are widely recognised along the south-central Andes (Chew *et al.*, 2007, 2008; Willner *et al.*, 2008; Bahlburg *et al.*, 2009). Indeed, the Mesoproterozoic and early Paleozoic-late Neoproterozoic detrital zircon components in particular are also commonly seen over thousands of kilometres, throughout East and West Antarctica and Australia, often in regions where possible source areas are surprisingly scarce or apparently absent. The examples along the South American Pacific margin cited above are usually within accretionary complexes where an autochthonous relationship at the Gondwana margin appears straightforward and probable. Comparing the U-Pb ages and Hf-isotope compositions of the detrital zircons from these complexes with possible Gondwanaland primary sources, the authors cited above all suggested sediment provenances in Paleozoic fold belts to the northeast and east, and more distantly in the Precambrian cratonic cores of Western Gondwanaland (southern Africa, East Antarctica, and central South America). There are abundant sources for the observed Mesoproterozoic and late Neoproterozoic zircons, admittedly by long (2,000 km) river transport, in southern Brazil, Uruguay, and northern Argentina. Closer sources of more limited outcrop, or known from drill-hole occurrences, of the required appropriate Paleozoic ages (Pankhurst *et al.*, 2003), were suggested in southern Argentina. However, the extensive Mesozoic-Cenozoic sedimentary cover there, and subsequent emplacement of voluminous Jurassic-Cretaceous igneous complexes in southern South America, makes the original extent and contemporary exposure of these

primary source areas very difficult to assess. The authors did not consider any western provenances as suggested in this work, and in any case, the important additional contribution from zircon Hf-isotope data is unavailable to assess its validity. Although a Zealandia provenance seems distant, it can be seen from figure 3, that these sources are no more distant than some alternatives in eastern South America, for example Brazil, Uruguay, or East Antarctica. Thus the well-developed Mesoproterozoic, early Paleozoic and Devonian-Carboniferous zircon age patterns in Chilean Patagonian metasedi-




FIG. 3. The Pacific margin of Gondwanaland in the late Paleozoic-Mesozoic showing the development of major contemporary depocentres at the Australia-Zealandia margin, and their subsequent, margin-parallel, dextral displacement as suspect terranes.

ments recommend further evaluation of a western, Zealandia, provenance.

The dominant Carboniferous, Permian and Triassic zircon proportions (>70%) in the Duque de York complex (Faúndez *et al.*, 2002; Lacassie *et al.*, 2006), and significant proportions (>20%) in the Chonos and Eastern Andean complexes (Hervé and Fanning, 2003; Hervé *et al.*, 2003) must indicate the proximity (although not seen) of contemporary volcanic arcs, nearer to the Patagonian margin of Gondwanaland, that effectively blocked the input of detrital zircons from older sources to the east. The nearest examples of such a magmatic arc would be in northern Patagonia (Augustsson *et al.*, 2006).

8. Conclusions

The basement of southern Zealandia was consolidated by Carboniferous times, and constituted extensive tracts latest Precambrian-Ordovician deep-marine turbiditic metasediments, intruded by Cambrian-Ordovician and Devonian-Carboniferous granitoids. A limited sedimentary cover comprised shallow-water Devonian rocks. There may have been local, but uncertain, extension of the Median Batholith (and several Permian-Triassic terranes) in New Zealand southeastwards along the Zealandia margin.

Substantial uplift of this basement progressed through late Paleozoic time (at least Silurian to Carboniferous), and the consequent erosion and river transport should have provided voluminous sediments, of mainly granitoid compositions, to postulated marine sedimentary basins at the Pacific margin of Zealandia. These sediments would have carried detrital zircon populations dominated by Mesoproterozoic, late Neoproterozoic-Cambrian-Ordovician, and Devonian-Carboniferous age components.

The present-day absence of any such late Paleozoic sedimentary basins at the southern Zealandia continental margin suggests that they may have been tectonically transported elsewhere during the Mesozoic, as suspect terranes in a dextral, margin-parallel sense, similar to that of the Eastern Province (Permian to Cretaceous) terranes of New Zealand. A possible destination for these terranes would be in the metamorphic complexes of Chilean Patagonia. Detrital zircon age populations in their metasediments compare well with those that might be expected in Zealandia

sedimentary basins, and thus a suspect terrane interpretation is permissible.

Acknowledgments

M. Fanning and co-workers are thanked for access to their primary Chilean zircon age datasets.

References

- Adams, C.J. 1981. Geochronological comparisons of Precambrian and Paleozoic orogens in New Zealand, Marie Byrd Land (West Antarctica), Northern Victoria Land (East Antarctica) and Tasmania. *In* Gondwana Five (Cresswell, M.M.; Vella, P.; editors). Selected papers and abstracts of papers presented at the Gondwana Symposium, No. 5: 191-192. Balkema, Rotterdam.
- Adams, C.J. 1986. Geochronological studies of the Swanson Formation of Marie Byrd Land, West Antarctica, and correlation with Northern Victoria Land, East Antarctica, and South Island, New Zealand. *New Zealand Journal of Geology and Geophysics* 29: 345-358.
- Adams, C.J. 1987. Geochronology of granite terranes in the Ford Ranges, Marie Byrd land, West Antarctica. *New Zealand Journal of Geology and Geophysics* 30: 57-72.
- Adams, C.J. 1997. Initial strontium isotopic signatures of Late Precambrian-Early Paleozoic metasediments from northern Victoria Land terranes, East Antarctic. *In* The Antarctic Region: Geological Evolution and Processes (Ricci, C-A.; editor). Proceedings of the International Symposium on Antarctic Earth Sciences, No. 7, Terra Antarctica Publication: 227-236. Siena, Italy.
- Adams, C.J. 2003. K-Ar geochronology of Torlesse Supergroup metasedimentary rocks in Canterbury, New Zealand. *Journal of the Royal Society of New Zealand* 33: 165-187.
- Adams, C.J. 2004. Rb-Sr age and strontium isotope characteristics of the Greenland Group, Buller terrane, New Zealand, and correlations at the East Gondwanaland margin. *New Zealand Journal of Geology and Geophysics* 47: 189-200.
- Adams, C.J. 2007. Paleozoic terranes at the Pacific Ocean margin of Zealandia Gondwana Research 13: 250-258.
- Adams, C.J.; Kreuzer, H. 1984. Potassium-argon ages of slates and phyllites from the Bowers and Robertson Bay Terranes, North Victoria Land, Antarctica. *In* German Antarctic North Victoria Land Expedition 1982/1983 GANOVEX III (Roland, N.; editor). *Geologisches Jahrbuch B6*: 265-288.

- Adams, C.J.; Kelley, S. 1998. Provenance of Permian-Triassic and Ordovician metagreywacke terranes in New Zealand: evidence from $^{40}\text{Ar}/^{39}\text{Ar}$ dating of detrital micas. *Geological Society of America Bulletin* 110: 422-432.
- Adams, C.J.; Maas, R. 2004. Age/isotopic characterisation of the Waipapa Group in Northland and Auckland, New Zealand, and implications for the status of the Waipapa Terrane. *New Zealand Journal of Geology and Geophysics* 47: 173-187.
- Adams, C.J.; Ireland, T.R. 2007. Provenance connections between Late Neoproterozoic and early Paleozoic sedimentary basins of the Ross Sea region, Antarctica, southeast Australia and southern Zealandia. In *Antarctic: A Keystone in a changing world* (Cooper, A.K.; Barrett, P.; Stagg, H.; Storey, B.; Stump, E.; Wise, W.; Davey, F.J.; Naish, T.R.; editors). Proceedings for the International Symposium on Antarctic Earth Sciences, No. 10. U.S. Geological Survey and the National Academies Open File Report 1047 Extended Abstracts:1-4.
- Adams, C.J.D.; Harper, C.T.; Laird, M.G. 1975. K-Ar ages of low grade metasediments of the Greenland and Waiuta Groups in Westland and Buller, New Zealand. *New Zealand Journal of Geology and Geophysics* 18: 39-48.
- Adams, C.J.; Morris, P.A.; Beggs, J.M. 1979. Age and correlation of volcanic rocks of Campbell Island and metamorphic basement of the Campbell Plateau, South-west Pacific. *New Zealand Journal of Geology and Geophysics* 6: 679-691.
- Adams, C.J.; Bishop, D.G.; Gabites, J.E. 1985. Potassium-argon age studies of a low-grade progressively metamorphosed greywacke sequence, Dansey Pass, South Island, New Zealand. *Journal of the Geological Society of London* 142: 339-349.
- Adams, C.J.; Barley, M.E.; Fletcher, I.R.; Pickard, A.L. 1998. Evidence from U-Pb zircon and $^{40}\text{Ar}/^{39}\text{Ar}$ muscovite detrital mineral ages in metasandstones for movement of the Torlesse suspect terrane around the eastern margin of Gondwanaland. *Terra Nova* 10: 183-189.
- Adams, C.J.; Pankhurst, R.J.; Maas, R.; Millar, I.L. 2005. Nd and Sr isotopic signatures of metasedimentary rocks around the South Pacific margin, and implications for their provenance. *Geological Society of London, Special Publication* 246: 113-142.
- Adams, C.J.; Campbell, H.J.; Griffin, W.J. 2007. Provenance comparisons of Permian to Jurassic tectonostratigraphic terranes in New Zealand: perspectives from detrital zircon age patterns. *Geological Magazine* 144: 701-729.
- Adams, C.J.; Campbell, H.J.; Griffin, W.L. 2009. Tracing the Caples Terrane through New Zealand using detrital zircon age patterns and radiogenic isotope signatures. *New Zealand Journal of Geology and Geophysics* 52: 223-245.
- Andrews, P.B. 1974. Deltaic sediments, Upper Triassic Torlesse Supergroup, Broken River, North Canterbury, New Zealand. *New Zealand Journal of Geology and Geophysics* 17: 881-905.
- Andrews, P.B.; Speden, I.G.; Bradshaw, J.D. 1976. Lithological and palaeontological content of the Carboniferous - Jurassic Canterbury Suite, South Island, New Zealand. *New Zealand Journal of Geology and Geophysics* 19: 791-819.
- Augustsson, C.; Münker, C.; Bahlburg, H.; Fanning, C.M. 2006. Provenance of late Palaeozoic metasediments of the SW South American Gondwana margin: a combined U-Pb and Hf-isotope study of single detrital zircons. *Journal of the Geological Society of London* 163: 983-995.
- Bahlburg, H.; Vervoort, J.D.; Du Frane, S.A.; Bock, B.; Augustsson, C.; Reimann, R.C. 2009. Timing of crust formation and recycling in accretionary orogens: Insights learned from the western margin of South America. *Earth-Science Reviews* 97: 227-253.
- Ballance, P.F.; Campbell, J.D. 1993. The Murihiku arc-related basin of New Zealand (Triassic-Jurassic). In *South Pacific Sedimentary Basins* (Ballance, P.F.; editor). *Sedimentary Basins of the World* 2: 21-33. Elsevier, Amsterdam.
- Birch, W.D. (editor) 2008 *Geology of Victoria*, Geological Society of Australia Special Publication 23, Chapter 6: 157-193.
- Bishop, G.J.; Bradshaw, J.D.; Landis, C.A. 1985. Provisional terrane map of South Island, New Zealand. In *Tectonostratigraphic Terranes, Circum-Pacific Council for Energy and Mineral resources* (Howell, D.G.; editor). *Earth Science Series* 1: 515-521. Houston, Texas.
- Bradshaw, J.D.; Andrews, P.B.; Field, B.D. 1983. Swanson Formation and related rocks of Marie Byrd Land and a comparison with the Robertson Bay Group of Northern Victoria Land. In *Antarctic Earth Science* (Oliver, R.L.; James, P.R.; Jago, J.B.; editors). *Australian Academy of Science*: 176-189. Canberra.
- Bradshaw, J.D.; Pankhurst, R.J.; Weaver, S.D.; Storey, B.C.; Muir, R.J.; Ireland, T.R. 1997. New Zealand superterrane recognized in Marie Byrd Land and Thurston Island. In *The Antarctic Region: Geological Evolution and Processes* (Ricci, C.-A.; editor). *Proceedings of the Seventh International Symposium*

- on Antarctic Earth Sciences, Siena, Terra Antarctica Publication: 429-436.
- Bradshaw, M.A. 1995. Stratigraphy and structure of the Lower Devonian rocks of the Waitahu and Orlando Outliers, near Reefton, New Zealand, and their relationship to the Inangahua Outlier. *New Zealand Journal of Geology and Geophysics* 38: 81-92.
- Bradshaw, M.A. 2000. Base of the Devonian Baton Formation and the question of a pre-Baton tectonic event in the Takaka terrane, New Zealand. *New Zealand Journal of Geology and Geophysics* 43: 601-610.
- Cawood, P.; Landis, C.; Nemchin, A.; Hada, S. 2002. Permian fragmentation, accretion and subsequent transition of a low-latitude Tehtyan sea-mount to the high latitude east Gondwana margin: evidence from detrital zircon age data. *Geological Magazine* 139: 131-144.
- Chew, D.M.; Schaltegger, U.; Oosler, J.; Whitehouse, M.J.; Gutjahr, M.; Spikings, R.A.; Miskovic, A. 2007. U-Pb geochronologic evidence for the evolution of the Gondwanan margin of the north-central Andes. *Geological Society of America Bulletin* 119: 697-711.
- Chew, D.M.; Magna, T.; Kirkland, C.L.; Miskovic, A.; Cardona, A.; Spikings, R.A.; Schaltegger, U. 2008. Detrital zircon fingerprint of the Proto-Andes: evidence for a Neoproterozoic active margin? *Precambrian Research* 167: 186-200.
- Coombs, D.S.; Landis, C.A.; Norris, R.J.; Sinton, J.M.; Borns, D.J.; Craw, D. 1976. The Dun Mountain Ophiolite Belt, New Zealand, its tectonic setting, constitution, and origin, with special reference to the southern portion. *American Journal of Science* 276: 561-603.
- Cooper, R.A.; Tulloch, A.J. 1992. Early Paleozoic terranes in New Zealand and their relationship to the Lachlan Fold Belt. *Tectonophysics* 214: 129-144.
- Faundez, V.; Hervé, F.; Lacassie, J.P. 2002. Provenance and depositional setting of pre-Late Jurassic turbidite complexes in Patagonia, Chile. *New Zealand Journal of Geology and Geophysics* 45: 411-426.
- Fioretti, A.M.; Black, L.P.; Foden, J.; Visona, D. 2005. Grenville age magmatism at the South Tasman Rise: a new piercing point for the reconstruction of Rodinia. *Geology* 33: 769-772.
- Foster, D.A.; Gray, D.R. 2000. Evolution and structure of the Lachlan Fold Belt (Orogen) of eastern Australia. *Annual Reviews of Earth and Planetary Sciences* 28: 47-80.
- Frost, C.D.; Coombs, D.S. 1989. Nd isotopic character of New Zealand sediments: implications for terrane concepts and crustal evolution. *American Journal of Science* 289: 744-770.
- Glen, R.G. 2005. The Tasmanides of eastern Australia. *In* *Terrane Processes at the margin of Gondwana* (Vaughan, A.P.M.; Leat, P.T.; Pankhurst, R.J.; editors). Geological Society of London Special Publication 246: 23-96.
- Gray, C.M. 1990. A strontium isotope traverse across the granitic rocks of southeastern Australia: Petrogenetic and tectonic implications. *Australian Journal of Earth Sciences* 37: 331-349.
- Grindley, G.W.; Mildenhall, D.C. 1981. Geological background to a Devonian plant fossil discovery, Ruppert Coast, Marie Byrd Land, West Antarctica. *In* *Gondwana Five* (Cresswell, M.M.; Vella, P.; editors): selected papers and abstracts of papers presented at the Gondwana Symposium, No. 5: 23-30. Balkema, Rotterdam.
- Hervé, F.; Fanning, C.M. 2003. Early Cretaceous subduction of continental crust at the Diego de Almagro archipelago, southern Chile. *Episodes* 26: 285-289.
- Hervé, F.; Fanning, C.M.; Pankhurst, R.J. 2003. Detrital zircon age patterns and provenance of the metamorphic complexes of southern Chile. *Journal of South American Earth Sciences* 16: 107-123.
- Ireland, T.R. 1992a. Correlation and provenance of sedimentary units around the southwestern Pacific margin. Geological Society of New Zealand, Annual Conference, Programme and Abstracts. Geological Society of New Zealand Miscellaneous Publication 63A: p. 84.
- Ireland, T.R. 1992b. Crustal evolution of New Zealand: Evidence from age distributions of detrital zircons in Western Province paragneisses and Torlesse greywacke. *Geochimica et Cosmochimica Acta* 56 (3): 911-920.
- Ireland, T.R.; Bradshaw, J.D.; Muir, R.; Weaver, S.D.; Adams, C. 1994. Zircon age distributions in granites, greywackes, and gneisses from the Southwest Pacific Gondwana region. *In* *Abstracts of the International Conference on Geochronology, Cosmochronology and Isotope Geology*, No. 8. U.S. Geological Survey, Circular 1107: p. 151.
- Ireland, T.R.; Flöttmann, T.; Fanning, C.M.; Gibson, G.M.; Preiss, W.V. 1998. Development of the early Palaeozoic Pacific margin of Gondwana from detrital-zircon ages across the Delamerian Orogen. *Geology* 26: 243-246.
- Jenkins, D.G.; Jenkins, T.B.H. 1971. First diagnostic Carboniferous fossils from New Zealand. *Nature* 233: 117-118.
- Kimbrough, D.L.; Tulloch, A.J.; Coombs, D.S.; Landis, C.A.; Johnston, M.R.; Mattinson, J.M. 1994. Uranium-

- lead zircon ages from the Median Tectonic Zone, South Island, New Zealand. *New Zealand Journal of Geology and Geophysics* 37: 393-419.
- Kreuzer, H.; Höhndorf, A.; Lenz, H.; Müller, P.; Vetter, U. 1987. Radiometric ages of Pre-Mesozoic rocks from Northern Victoria Land, Antarctica. *In* *Gondwana Six: Structure, Tectonics and Geophysics* (McKenzie, G.D.; editor). American Geophysical Union Monograph 40: 31-48.
- Lacassie, J.P.; Hervé, F.; Roser, B. 2006. Sedimentary provenance of the post-Early Permian to pre-Early Cretaceous metasedimentary Duque de York Complex, Chile. *Revista Geológica Chile* 33 (2): 199-219.
- Landis, C.A.; Campbell, H.J.; Aslund, T.; Cawood, P.A.; Douglas, A.; Kimbrough, D.L.; Pillai, D.D.L.; Raine, J.I.; Willsman, A. 1999. Permian-Jurassic strata at Productus Creek, Southland, New Zealand: implications for terrane dynamics at the eastern Gondwanaland margin. *New Zealand Journal of Geology and Geophysics* 42: 255-278.
- MacKinnon, T.C. 1983. Origin of the Torlesse Terrane and coeval rocks. *Geological Society of America Bulletin* 94: 967-985.
- Mortimer, N.; Tulloch, A.J.; Gans, P.; Calvert, A.; Walker, N. 1999. Geology and thermochronometry of the east edge of the Median Tectonic Batholith (Median Tectonic Zone): a new perspective on Permian to Cretaceous crustal growth of New Zealand. *The Island Arc* 8: 404-425.
- Mortimer, N.; Davey, F.J.; Melhuish, A.; Yu, J.; Godfrey, N.J. 2002. Geological interpretation of a deep seismic reflection profile across the Eastern province and Median Batholith, New Zealand: crustal architecture of an extended Phanerozoic convergent orogen. *New Zealand Journal of Geology and Geophysics* 45: 349-364.
- Muir, R.J.; Ireland, T.R.; Weaver, S.D.; Bradshaw, J.D. 1996. Ion microprobe dating of Paleozoic granitoids: Devonian magmatism in New Zealand and correlations with Australia and Antarctica. *Chemical Geology (Isotope Geosciences Section)* 127: 191-210.
- Pankhurst, R.J.; Weaver, S.D.; Bradshaw, J.D.; Storey, B.C.; Ireland, T.R. 1998. Geochronology and geochemistry of pre-Jurassic superterrane in Marie Byrd land, Antarctica. *Journal of Geophysical Research B* 103: 2529-2547.
- Pankhurst, R.J.; Rapela, C.W.; Loske, W.P.; Márquez, M.; Fanning, C.M. 2003. Chronological study of the pre-Permian rocks of southern Patagonia. *Journal of South American Earth Sciences* 16: 22-41.
- Pickard, A.L.; Adams, C.J.; Barley, M.E. 2000. Australian provenances for Upper Permian to Cretaceous rocks forming accretionary complexes on the New Zealand sector of the Gondwanaland margin. *Australian Journal of Earth Sciences* 47: 987-1007.
- Pickett, D.A.; Wasserburg, G.J. 1989. Neodymium and strontium isotopic characteristics of New Zealand granitoids and related rocks. *Contributions to Mineralogy and Petrology*, 103: 131-142.
- Rattenbury, M.S.; Cooper, R.A.; Johnston, M.R. (compilers). 1998. *Geology of the Nelson Area*. Institute of Geological and Nuclear Sciences, Geological map 9, scale 1:250,000: 67 p. Lower Hutt, New Zealand.
- Retallack, G.J. 1983. Middle Triassic estuarine deposits near Benmore Dam, southern Canterbury and northern Otago. *Journal of the Royal Society of New Zealand* 13: 107-127.
- Retallack, G.J.; Ryburn, R.J. 1982. Middle Triassic deltaic deposits in Long Gully, near Otematata, north Otago, New Zealand. *Journal of the Royal Society of New Zealand* 12: 207-227.
- Roser, B.P.; Korsch, R.J. 1999. Geochemical characterisation, evolution and source of a Mesozoic accretionary wedge, the Torlesse terrane, New Zealand. *Geological Magazine* 136: 493-512.
- Smale, D. 1980. Akatarawa Conglomerate (Permian), Lake Aviemore, South Canterbury. *New Zealand Journal of Geology and Geophysics* 23: 279-292.
- Sutherland, R. 1999. Basement geology and tectonic development of the greater New Zealand region: an interpretation from regional magnetic data. *Tectonophysics* 308: 341-362.
- Tessensohn, F.; Duphorn, K.; Jordan, H.; Kleinschmidt, G.; Skinner, D.N.B.; Vetter, U.; Wright, T.O.; Wyborn, D. 1981. Geological comparison of basement units in North Victoria Land, Antarctica. *In* *Ganovex German Antarctic North Victoria Expedition 1979/80*. *Geologisches Jahrbuch B41*: 31-88.
- Tonarini, S.; Rocchi, S. 1994. Geochronology of Cambro-Ordovician intrusives in Northern Victoria Land: A review. *Terra Antarctica* 1: 59-62.
- VandenBerg, A.H.M.; Willman, C.E.; Maher, S.; Cayley, R.E.; Taylor, D.H.; Morand, M.J.; Simons, B.A.; Moore, D.H.; Radojkovic, A. 2000. *The Tasman Fold Belt System in Victoria*. Geological Survey of Victoria Special Publication.
- Vita, G.; Lombardo, B.; Guiliani, O. 1991. Ordovician uplift pattern in the Wilson Terrane of the Borch-grevink Coast, Victoria Land, Antarctica. *Proceedings of the meeting Earth Science investigations in Antarctica*,

- Siena, 4-6 October 1989. *Memoria Societa Italiana* 46: 257-265.
- Veevers, J.J. (editor) 2000. Billion-year history of Australia and neighbours in Gondwanaland. Gemoc Press: 388 p.
- Wandres, A.M.; Bradshaw, J.D.; Weaver, S.; Maas, R.; Ireland, T.R.; Eby, N. 2004a. Provenance of the sedimentary Rakaia subterrane, South Island, New Zealand: the use of igneous clast compositions to define the source. *Sedimentology* 168: 193-226.
- Wandres, A.M.; Bradshaw, J.D.; Weaver, S.; Maas, R.; Ireland, T.R.; Eby, N. 2004b. Provenance analysis using conglomerate clast lithologies: a case study from the Pahau terrane of New Zealand. *Sedimentology* 167: 57-89.
- Willner, A.P.; Gerdes, A.; Masonne, H.J. 2008. History of crustal growth and recycling at the Pacific convergent margin of South America at latitudes 29°-36° revealed by a U-Pb and Lu-Hf isotope study of detrital zircon from late Paleozoic accretionary systems. *Chemical Geology* 253: 114-129.
- Wright, T.O.; Dallmeyer, R.D. 1991. The age of cleavage development in the Ross orogen, northern Victoria Land, Antarctica: evidence from ⁴⁰Ar/³⁹Ar whole-rock slate ages. *Journal of Structural Geology* 13: 677-690.